New Initiators of Opioid Use Disorder (OUD) Treatment with MAT, Quarterly

Updated: August 29, 2019

Metric Information

Metric description: Percent (%) of Medicaid enrollees with a history of OUD, untreated in the prior calendar quarter, treated by any of the various medication assisted treatment (MAT) modalities for OUD in a given calendar quarter. A single rate is reported.

Metric specification version: Developed by ARM/HCA with Assistance from DSHS/RDA

Data collection method: Administrative only.

Data source: ProviderOne Medicaid claims/encounter and enrollment data.

Claim status: Include only final paid claims or accepted encounters in Metric calculation.

Identification window: Measurement Quarter for numerator and 2-year window from Measurement Quarter for denominator.

Direction of quality improvement: Higher is better.

Metric Details

Eligible Population		
Age	N/A	
Gender	N/A	
Minimum Medicaid	Month of service when the diagnosis was made, OUD treatment	
enrollment	provided	
Allowable gap in Medicaid	N/A	
enrollment		
Medicaid enrollment anchor	Last day of the Measurement Quarter	
date		
Medicaid benefit and	Includes Medicaid beneficiaries with comprehensive medical	
eligibility	benefits. Excludes beneficiaries that are eligible for both Medicare	
	and Medicaid, beneficiaries with primary insurance other than	
	Medicaid.	

Denominator:

Data elements required for denominator: Medicaid beneficiaries who meet the above eligibility criteria are identified and following steps are needed to identify the eligible Medicaid beneficiaries for the denominator.

- Determine the last day of the reporting quarter.
- In claim history going back 2 years for those that qualify in the reporting quarter, identify those with any diagnosis of OUD using the DSHS/RDA OUD Diagnosis Narrow Value Set as a diagnosis in any setting

New Initiators of Opioid Use Disorder (OUD) Treatment with MAT, Quarterly

- The qualified individuals with diagnosis serve as a denominator for the reporting calendar quarter

Required exclusions for denominator.

- Eligible population exclusions:
 - o Beneficiaries that are eligible for both Medicare and Medicaid.
 - o Beneficiaries with primary insurance other than Medicaid.
 - o Beneficiaries with no diagnosis of OUD Narrow in past 2 year period

Value sets required for denominator.

Name	Value Set
OUD Diagnosis Narrow Value Set	See Attached from DSHS/RDA

Numerator:

Beneficiaries must qualify for inclusion in the denominator to be eligible for inclusion in the numerator.

Data elements required for numerator: Identification of paid claims during the reporting quarter for individuals that meet the following criteria.

- (1) Any pharmacy claim within reporting quarter, with day supply > 0, for drugs identified by GPIs under MAT_type of 'BupNaloxoneCombo' or, 'Naltrexone' or 'Buprenorphine' in the OUD Treatment MAT Value Set or,
 - Any paid claim in any setting, within reporting quarter, for procedure codes that meet treatment with 'Opiate Substitution' from the OUD Treatment Opiate Substitution Value Set
- (2) No pharmacy claim in the prior quarter, for any drugs identified by GPIs under MAT_type of 'BupNaloxoneCombo' or, 'Naltrexone' or 'Buprenorphine' in the OUD Treatment MAT Value Set and,
 - no paid claim in any setting, in the prior quarter, with procedure codes that meet treatment with 'Opiate Substitution' from the OUD Treatment Opiate Substitution Value Set
- (3) Count the number of unique beneficiaries after step 2 to calculate the numerator.

Measure Rate:

Divide the numerator by denominator for rate

Value sets required for numerator.

Name	Value Set
OUD Treatment Opiate Substitution Value Set	See Attached from DSHS/RDA
OUD Diagnosis Narrow Value Set	See Attached from DSHS/RDA
OUD Treatment MAT Value Set	See Attached from DSHS/RDA

Required exclusions for numerator.

- None

Deviations from cited specifications for numerator.

- None

New Initiators of Opioid Use Disorder (OUD) Treatment with MAT, Quarterly

Version Control

December 2018, v1.1

August 2019, v1.2 Updates include limiting eligible population with a narrow diagnosis of OUD